

*Stenocarus umbrinus* (Gyllenhal) — 6.vii.1980. Usually associated with field edges or disturbed ground, rather than fenland, it is likely that the single example found at Pashford was a 'stray' from an adjacent cornfield, red with Poppies, *Papaver rhoeas* L., with which the species is associated.

*Ceutorhynchus angulosus* Boheman — 15.vi.1980. Since this was recorded 'new to Suffolk' (Mendel, H., 1983, *Trans. Suffolk Nat. Soc.*, 19: 337-40) I have been informed that Dr. M. G. Morris also took a specimen at Pashford, on 18th. June 1969 (Morris, *in litt.*). *Acalyptus carpini* (F.) — 10.viii.1980 & 22.v.1981. First recorded from Pashford Poors Fen, and Suffolk, on 10th August 1980 and shortly afterwards at the same site by Dr. R. C. Welch on 26th. August 1980 (Suffolk Biological Records Centre, unpublished ms.), this fenland species is associated with Sallows, *Salix* spp. *A. carpini* is well known from Wicken Fen, Cambridgeshire, and has been recorded from Great Fen, Lopham, just over the Suffolk border in Norfolk (Pope, R. D., 1968, *Trans. Suffolk Nat. Soc.*, 14: 25-40).

I thank the Suffolk Trust for Nature Conservation for permission to collect at Pashford Poors Fen Nature Reserve, Dr. M. L. Cox for identifying *Longitarsus brunneus*, Drs. M. G. Morris and R. C. Welch, and Messrs. J. W. Digby, E. Milne-Redhead and D. R. Nash for allowing me to include their unpublished records. — H. MENDEL, The Museum, High Street, Ipswich, Suffolk IP1 3QH.

TWO ADDITIONAL SITES FOR EUDECTUS WHITEI SHARP (COL.: STAPHYLINIDAE) IN SCOTLAND. — The paucity of recorded sites for *E. whitei* in Britain prompts me to record somewhat belatedly that I took a single example from fine gravel among sparse vegetation on an exposed ridge on Sgurr Mhor, Wester Ross (1110m) about 100m below the summit in July 1982 and two examples on Ben Lawers, Perthshire (1214m) on the south side just below the summit in May 1984. These would appear to be respectively the most northerly and most southerly records for the species in Scotland though it is, of course, recorded from a few sites in the Penines. J. A. OWEN 8 Kingsdown Road, Epsom, Surrey, KT17 3PU.

OTIORHYNCHUS PORCATUS (HERBST) (COL.: CURCULIONIDAE) IN LEICESTERSHIRE — Pitfall traps containing a 5 per cent solution of formalin have been operated in my garden in Leicester since 1979 starting each year on 1 April and ending on 31 October. All beetles trapped in 1979-84 have been identified by Dr. D. Goddard. Among them are 31 *Otiorhynchus porcatus*, distributed as follows: 1979 (2), 1980 (2), 1981 (4), 1982 (5), 1983 (14) and 1984 (4). This suggests an established population of a species which is new to Leicestershire. DENIS F. OWEN, 66 Scraftoft Lane, Leicester LE5 1HU.